

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Great Plains Wildlife Damage Control Workshop Wildlife Damage Management, Internet Center
Proceedings for

12-1-1979

IMPACT OF PLAINS POCKET GOPHERS ON FORAGE PRODUCTION

Ronald M. Case

University of Nebraska-Lincoln, rcase2@neb.rr.com

James L. Stubbendieck

University of Nebraska-Lincoln, jstubbendieck@unl.edu

Dale G. Luce

Nebraska Public Power District, Columbus, Nebraska

Follow this and additional works at: <https://digitalcommons.unl.edu/gpwdcwp>

 Part of the [Environmental Health and Protection Commons](#)

Case, Ronald M.; Stubbendieck, James L.; and Luce, Dale G., "IMPACT OF PLAINS POCKET GOPHERS ON FORAGE PRODUCTION" (1979). *Great Plains Wildlife Damage Control Workshop Proceedings*. 240.
<https://digitalcommons.unl.edu/gpwdcwp/240>

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Great Plains Wildlife Damage Control Workshop Proceedings by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

IMPACT OF PLAINS POCKET GOPHERS ON FORAGE PRODUCTION

Ronald M. Case, Department of Forestry, Fisheries and Wildlife, University of Nebraska, James L. Stubbendieck, Department of Agronomy, University of Nebraska, Lincoln 68583 and Dale G. **Luce**, Nebraska Public Power District, Columbus, Nebraska 68601

The presence of pocket gophers on rangeland and farmland is highly visible due to the earth mounds which they build. The nuisance impact of their mounds and acknowledged long-term beneficial effects on soil (**Laycock** and Richardson 1975) result in a value which is debatable. The impact of various species of Thomomys on **herbage** production of rangelands has been widely reported (Fitch and Bentley 1949; **Richens** 1965; Turner 1969; **Laycock** and Richardson 1975; **Alsager** 1977). However, besides our studies in Nebraska, we could find no literature concerning the effect of Geomys on forage production and no literature concerning the effect of any gopher on alfalfa production.

The objective of our paper is to review the impact of plains pocket gophers (Geomys bursarius) on forage production. This includes the effects of gophers on alfalfa production.

METHODS

Rangeland. This study area is in Dawes County in the panhandle of Nebraska, about 56 km southeast of **Chadron**. Four range sites on the study area were selected. Two 0.64 ha areas were selected on each site, one with gophers present and one without. Each of the 0.64 ha areas was subdivided into four replicates. In September of 1975 and 1976 the areas were harvested to a height of 2.5 cm above ground level. Each range site also had a cattle **exclosure** which was harvested at the same time. Harvest data were adjusted upward to ungrazed status and all weight data are on an oven-dry basis.

Alfalfa Field. This study area is in Lancaster County, Nebraska, approximately 8 km east of Lincoln. Two alfalfa fields were selected. On each field four paired areas were chosen, one of each pair had gophers present while they were absent on the other. Each area was harvested to a height of 2.5 cm above ground level just prior to the landowner's harvesting of the fields during 1978. Mean oven-dry weights of three cuttings were computed.

RESULTS

Range condition within each of the four sites did not differ during the two year study. The condition of the four sites were different from each other, however, and the areas where gophers were present were usually lower and never higher in range condition than the areas where they were absent (Table 1).

Table 1. Range condition (Nichols et al. 1978) on four sites in western Nebraska with and without plains pocket gophers.

Range	Site	Gophers Present	Range Condition	
	Sands	Yes	Good	
		No	High	Good
	Sands	Yes	Good	
		No	Excellent	
	Silty	Yes	Low	Fair
		No	High	Fair
	Silty	Yes	Low	Fair
		No	Low	Fair

Forage yields on each of the range sites were significantly less ($P < 0.05$) on areas where gophers were present compared to where they were absent (Table 2). The 2-year average decrease in forage production due to the presence of gophers ranged from 21% to 49% on the different sites.

Table 2. Effect of plains pocket gophers on forage yield on four range sites in western Nebraska.

Range	Site	Gophers Present	Year		Average	Decrease
			1975	1976		
			kg/ha			
	Sands	Yes	1054	1411	1233	36%
		No	1669	2155	1912	
	Sands	Yes	1105	1801	1453	47%
		No	2107	3338	2723	
	Silty	Yes	665	672	669	49%
		No	1307	1316	1312	
	Silty	Yes	638	605	622	21%
		No	778	806	792	

Gophers reduced forage yield on alfalfa fields by an amount similar to that on rangeland (Table 3). The yield of alfalfa was significantly less ($P < 0.05$) and the weed yield significantly higher ($P < 0.05$) opportunities of the field where gophers were present compared to areas where they were absent. Although the weed yield increased dramatically, the combined yield of alfalfa and weeds where gophers were present was 38% less than where gophers were absent.

Table 3. Effect of plains pocket gophers on alfalfa and weed yield in alfalfa fields in southeastern Nebraska.

Gophers 'Present	Alfalfa Yield	Weed Yield kg /ha	Combined Yield	Decrease
Yes	5577	741	6318	38%
No	10017	105	10122	

DISCUSSION

In addition to decreased forage yields, plains pocket gophers increased the percentage of bare soil, decreased basal cover and decreased perennial grasses with a concomitant increase in annual grasses on the rangeland study area in western Nebraska (Foster and Stubbendieck 1980). Alfalfa plant density was decreased and frequency of weed species occurrence increased on gopher inhabited areas of alfalfa fields (Luce 1979).

When examining the significance of gophers on yield and considering possible control methods, the following is important: if gophers decrease yield 50% but occupy only 10% of the area, then the total loss in forage production is only 5%. Control decisions are then based upon the concentration of the gopher population and time and money necessary for control.

Pocket gophers are often thought to be characteristic of overgrazed land (Phillips 1936). In fact, Mielke (1977) suggested that gophers complemented the overgrazing and trampling of bison which resulted in increased forbs. The gophers, he suggested, then thrived and in working the soil stimulated vegetation growth suitable for bison. Myers and Vaughan (1964) and Luce et al. (1980) both documented that plains pocket gophers eat primarily grasses. When examining range condition in this study, it appears that the activity of gophers sets back succession. Thus the poorer range conditions may not be a prerequisite for gopher habitation; the gophers are likely a cause of poorer range conditions.

ACKNOWLEDGEMENTS

Thanks to Drs. R. Johnson, E. Peters and R. Timm for their critical review of this manuscript. J. Andelt diligently typed several drafts and the final paper.

LITERATURE CITED

- ALSAGER, D.E. 1977. Impact of pocket gophers (Thomomys talpoides) on the quantitative productivity of rangeland vegetation in southern Alberta: a damage assessment tool. Pages 47-57 in W.B. Jackson and R.E. Marsh, eds., Test Methods for Vertebrate Pest Control and Management Materials, ASTM STP 625, American Society for Testing and Materials. 256pp.
- FITCH, H.S., and J.R. BENTLEY. 1949. Use of California annual-plant forage by range rodents. Ecology 30:306-321.
- FOSTER, M.A., and J. STUBBENDIECK. 1980. Effects of the plains pocket gopher (Geomys bursarius) on rangeland. J. Range Manage. 31:74-78.
- LAYCOCK, W.A., and B.A. RICHARDSON. 1975. Long-term effects of pocket gopher control on vegetation and soils of a subalpine grassland. J. Range Manage. 28:458-462.
- LUCE, D.G. 1979. Plains pocket gopher (Geomys bursarius) damage to alfalfa. M.S. Thesis. Univ. Nebraska, Lincoln. 63pp.
- LUCE, D.G., R.M. CASE, and J. STUBBENDIECK. 1980. Food habits of the plains pocket gopher on western Nebraska rangeland. J. Range Manage. In Press.
- MIELKE, H.W. 1977. Mound building by pocket gophers (Geomyidae): Their impact on soils and vegetation in North America. J. Biogeography 4:171-180.
- MYERS, G.T., and T.A. VAUGHAN. 1964. Food habits of the plains pocket gopher in eastern Colorado. J. Mammal. 45:588-598.
- NICHOLS, J.T., P.N. JENSEN, and J. STUBBENDIECK. 1978. Range judging handbook for Nebraska. University of Nebraska Cooperative Extension Service, Extension Circular 1-37-78. 30pp.
- PHILLIPS, P. 1936. The distribution of rodents in overgrazed and normal grasslands of central Oklahoma. Ecology 17:673-679.
- RICHERS, V.B. 1965. An evaluation of control on the Wasatch pocket gopher. J. Wildlife Manage. 29:413-425.
- TURNER, G.T. 1969. Responses of mountain grassland vegetation to gopher control, reduced grazing, and herbicide. J. Range Manage. 22:377-383.